Stanford Exploration Project, Report 79, November 17, 1997, pages 17–22

Short Note

3-D Eikonal equation for time migration

Arnaud Berlioux[1]

keywords: *three-dimensional, time migration, velocity*

INTRODUCTION

In the short-note I wrote in the previous SEP report (Berlioux, 1993b), I presented a new method to perform depth migration in 3-D, starting from time-migrated maps. Since time migration does not always produce properly migrated images or maps in areas where velocity variations are present (Hatton et al., 1981), the method I proposed consists of two parts: first undo the time migration by a principle of *de-migration*, and then perform the depth migration.

In my previous work these two parts were based on a ray-tracing algorithm, and the de-migration process corresponded to the inverse operation of Kirchhoff time migration. De-migration and time migration have to remain associated as inverse processes. I now investigate the method introduced in 1991 by V.J. Khare to perform time migration in order to produce a de-migration scheme corresponding to this process.

In this paper, I explain briefly how V.J. Khare derived the 2-D Eikonal equation for time migration and what modifications have to be introduced to handle the 3-D case, and I make a proposal to use this equation for the de-migration process.

INTRODUCTION OF EIKONAL EQUATION FOR TIME-MIGRATION

In two dimensions, considering one-way time, the wave equation is

$$\frac{\partial^2 p}{\partial x^2} + \frac{\partial^2 p}{\partial z^2} = \frac{1}{V^2(x,z)} \frac{\partial^2 p}{\partial t^2} \tag{1}$$

where $p(x,z,t)$ is the pressure wave field, and $V(x,z)$ is the velocity of the wave propagation in the medium.

[1]**email:** not available

We can make the following approximation of the wave field:

$$p\,(x,z,t) \;=\; \hat{p}\,(x,z)\,e^{\,i\,\omega\,(t\,-\,\tau\,(x,z)\,)} \tag{2}$$

expressing the pressure wave field in term of amplitude $\hat{p}(x,z)$ and phase $\omega\,(t\,-\,\tau\,(x,z)\,)$, where τ is the traveltime along the ray paths. Then replacing (2) in equation (1) and making the high frequency approximation, we obtain the 2-D Eikonal equation:

$$\left(\frac{\partial\tau}{\partial x}\right)^2 \;+\; \left(\frac{\partial\tau}{\partial z}\right)^2 \;=\; \frac{1}{V^2\,(x,z)} \tag{3}$$

Among other uses, this equation is useful to compute traveltimes of rays in a given earth model.

V.J. Khare introduced the following equation for the ray-theoretic interpretation of *all-angle* time migration (Khare, 1991):

$$\frac{\partial^2 p'}{\partial x'^2} \;+\; \frac{1}{\tilde{V}^2\,(x',\zeta)}\,\frac{\partial^2 p'}{\partial\zeta^2} \;=\; -\,\frac{2}{\tilde{V}^2(x',\zeta)}\,\frac{\partial^2 p'}{\partial t'\,\partial\zeta} \tag{4}$$

where $p'\,(x',\zeta,t')$ and $\tilde{V}\,(x',\zeta)$ are respectively the pressure wave field and the wave propagation velocity in the time-retarded coordinate system (Claerbout, 1976),

$$\begin{aligned} x' &= x \\ t' &= t \,+\, \zeta \end{aligned} \tag{5}$$

with the time-like vertical coordinate ζ (Lowenthal et al., 1985),

$$\zeta \;=\; \int_0^z \frac{dz}{V\,(x,z)} \tag{6}$$

If we make the following approximation of the wave field in the new coordinate system, equivalent to the equation (2):

$$p'\,(x',\zeta,t') \;=\; \hat{p'}\,(x',\zeta)\,e^{\,i\,\omega\,(t'\,-\,\tau'\,(x',\zeta)\,)} \tag{7}$$

we can express the 2-D Eikonal equation for time migration giving the traveltime τ' in the frequency domain in the time-retarded coordinate system:

$$\left(\frac{\partial\tau'}{\partial x'}\right)^2 \;+\; \frac{1}{\tilde{V}^2\,(x',\zeta)}\left(\frac{\partial\tau'}{\partial\zeta}\right)^2 \;=\; \frac{2}{\tilde{V}^2\,(x',\zeta)}\,\frac{\partial\tau'}{\partial\zeta} \tag{8}$$

Using equations (5) and (6), we obtain the 2-D Eikonal equation for time migration in the physical coordinate system:

$$\left(\frac{\partial\tau}{\partial x} \;+\; \mathcal{A}\,(x,z)\,\frac{\partial\tau}{\partial z}\right)^2 \;+\; \left(\frac{\partial\tau}{\partial z}\right)^2 \;=\; \frac{1}{V^2\,(x,z)} \tag{9}$$

with

$$\mathcal{A}\,(x,z) \;=\; V\,(x,z)\,\int_0^z \frac{V_x\,(x,z)}{V^2\,(x,z)}\,dz \;=\; -\,V\,(x,z)\,\frac{\partial\zeta}{\partial x} \tag{10}$$

USING 3-D EIKONAL EQUATION FOR DE-MIGRATION

Presentation of the 3-D case

In the 3-D case, the wave equation corresponding to the ray-theoretic interpretation of *all-angle* time migration is

$$\frac{\partial^2 p'}{\partial x'^2} + \frac{\partial^2 p'}{\partial y'^2} + \frac{1}{\tilde{V}^2\left(x', y', \zeta\right)} \frac{\partial^2 p'}{\partial \zeta^2} = -\frac{2}{\tilde{V}^2(x', y', \zeta)} \frac{\partial^2 p'}{\partial t' \, \partial \zeta} \tag{11}$$

in the 3-D time-retarded coordinate system

$$\begin{aligned} x' &= x \\ y' &= y \\ t' &= t + \zeta \end{aligned} \tag{12}$$

with the new time-like vertical coordinate ζ,

$$\zeta = \int_0^z \frac{dz}{V\left(x, y, z\right)} \tag{13}$$

Following the same scheme as for the 2-D case and using equations (12) and (13), we finally obtain the 3-D Eikonal equation for time migration:

$$\left(\frac{\partial \tau}{\partial x} + \mathcal{A}\left(x, y, z\right) \frac{\partial \tau}{\partial z}\right)^2 + \left(\frac{\partial \tau}{\partial y} + \mathcal{B}\left(x, y, z\right) \frac{\partial \tau}{\partial z}\right)^2 + \left(\frac{\partial \tau}{\partial z}\right)^2 = \frac{1}{V^2\left(x, y, z\right)} \tag{14}$$

with

$$\mathcal{A}\left(x, y, z\right) = -V\left(x, y, z\right) \frac{\partial \zeta}{\partial x} \tag{15}$$

and

$$\mathcal{B}\left(x, y, z\right) = -V\left(x, y, z\right) \frac{\partial \zeta}{\partial y} \tag{16}$$

Using the Eikonal equation for de-migration: a proposal

It is obvious that this equation in 3-D cannot be solved analytically for any kind of velocity model. Since the Eikonal equation relates the gradient of the traveltime τ to the velocity structure, several methods have been formulated to study the classical Eikonal equation (3) for a velocity structure sampled at discrete points in a 3-D space (3-D grid) and using finite-difference algorithms: M. Reshef and D. Kosloff (1986) integrate a finite-difference approximation of the Eikonal equation with a Runge-Kutta method; J. Vidale (1988) also uses a finite-difference algorithm but solves directly for traveltimes using a planar or circular wavefront extrapolation; and J. Van Trier and W. Symes (1990) derive a conservation law from the Eikonal equation depicting the first arrival time field.

Each differential term in Eikonal equation (14) can be approximated with finite difference to give the traveltime at a particular point of a grid with respect to the previously calculated traveltimes. To apply a finite-difference scheme to this equation, we need a 3-D grid of velocities, and since I use GOCAD to handle 3-D surfaces and structures, I present in this report a method to compute a 3-D grid of velocities from a GOCAD model (Berlioux, 1993a).

The aim of this project is to find how we can obtain a de-migration process from this Eikonal equation for time-migration. Nevertheless, to find an "inverse" process to the time migration method presented here, it is necessary to understand clearly how it is performed. I will therefore try to operate the time migration on several synthetic models of zero-offset data.

CONCLUSIONS

Until now, I have not tried to implement any of the methods proposed in the papers mentioned above. Most of them call for finite-difference algorithms, and it appears to me necessary to find a way to produce 3-D grids easily and rapidly from a 3-D model.

The Eikonal equation for time migration introduced by V.J. Khare and the theory that comes with it seem to represent an interesting way to obtain a de-migration process accurate enough to handle most of the complicated models.

ACKNOWLEDGMENTS

I would like to thank Mr. V.J. Khare for having devoted part of his time to me.

REFERENCES

Berlioux, A., 1993a, 3-D grid with GOCAD: SEP–**79**, 301–318.

Berlioux, A., 1993b, Three-dimensional map de-migration/migration: SEP–**77**, 71–76.

Claerbout, J. F., 1976, Fundamentals of geophysical data processing: Blackwell.

Hatton, L., Larner, K. L., and Gibson, B. S., 1981, Migration of seismic data from inhomogeneous media: Geophysics, **46**, no. 5, 751–767.

Khare, V., 1991, A new ray-theoretic interpretation of time migration: 61st Annual Internat. Mtg., Soc. Expl. Geophys., Expanded Abstracts, 1492–1496.

Lowenthal, D., Lu, L., Roberson, R., and Sherwood, J. W. C., 1985, The wave equation applied to migration *in* Gardner, G. H. F., Ed., Migration of seismic data:: Soc. Expl. Geophys., 208–227.

Reshef, M., and Kosloff, D., 1986, Migration of common shot gathers: Geophysics, **51**, no. 2, 324–331.

van Trier, J., and Symes, W. W., 1990, Upwind finite difference calculation of traveltimes: SEP–**65**, 41–58.

Vidale, J., 1988, Finite-difference calculation of travel times: Bull. Seism. Soc. Am., **78**, 2062–2076.